

Succinic acid, 4-chloro-2-nitrobenzyl pentyl ester

| | |
|----------------------|--|
| Inchi: | InChI=1S/C16H20ClNO6/c1-2-3-4-9-23-15(19)7-8-16(20)24-11-12-5-6-13(17)10-14(12)1 |
| InchiKey: | WWGSJZFAHHLUTK-UHFFFAOYSA-N |
| Formula: | C16H20ClNO6 |
| SMILES: | CCCCCOC(=O)CCC(=O)OCc1ccc(Cl)cc1[N+](=O)[O-] |
| Mol. weight [g/mol]: | 357.79 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|---------|----------------------|----------------|
| gf | -267.23 | kJ/mol | Joback Method |
| hf | -676.08 | kJ/mol | Joback Method |
| hfus | 51.59 | kJ/mol | Joback Method |
| hvap | 94.10 | kJ/mol | Joback Method |
| log10ws | -5.18 | | Crippen Method |
| logp | 3.805 | | Crippen Method |
| mcvol | 257.080 | ml/mol | McGowan Method |
| pc | 1756.54 | kPa | Joback Method |
| rinpol | 2542.00 | | NIST Webbook |
| rinpol | 2542.00 | | NIST Webbook |
| tb | 943.97 | K | Joback Method |
| tc | 1171.44 | K | Joback Method |
| tf | 639.39 | K | Joback Method |
| vc | 1.002 | m ³ /kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|--------|---------|-----------------|---------------|
| cpg | 774.31 | J/molxK | 943.97 | Joback Method |
| cpg | 785.02 | J/molxK | 981.88 | Joback Method |
| cpg | 794.53 | J/molxK | 1019.79 | Joback Method |
| cpg | 802.86 | J/molxK | 1057.70 | Joback Method |
| cpg | 810.02 | J/molxK | 1095.61 | Joback Method |
| cpg | 816.03 | J/molxK | 1133.52 | Joback Method |
| cpg | 820.92 | J/molxK | 1171.44 | Joback Method |

Sources

| | |
|------------------------|---|
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=U380938&Units=SI |
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci990307l |
| Crippen Method: | https://www.cheméo.com/doc/models/crippen_log10ws |

Legend

| | |
|----------------------------|---|
| cp_g: | Ideal gas heat capacity |
| g_f: | Standard Gibbs free energy of formation |
| h_f: | Enthalpy of formation at standard conditions |
| h_{fus}: | Enthalpy of fusion at standard conditions |
| h_{vap}: | Enthalpy of vaporization at standard conditions |
| log₁₀ws: | Log ₁₀ of Water solubility in mol/l |
| log_p: | Octanol/Water partition coefficient |
| mc_{vol}: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rin_{pol}: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.cheméo.com/cid/120-884-7/Succinic-acid-4-chloro-2-nitrobenzyl-pentyl-ester.pdf>

Generated by Cheméo on 2024-05-01 23:16:49.30417888 +0000 UTC m=+16894658.224756197.

Cheméo (<https://www.cheméo.com>) is the biggest free database of chemical and physical data for the process industry.