1H-Imidazole, 1-methyl-

Other names:	1-Methyl-1H-imidazole
	1-Methylimidazole
	Imidazole, 1-methyl-
	N-Methylimidazole
Inchi:	InChI=1S/C4H6N2/c1-6-3-2-5-4-6/h2-4H,1H3
InchiKey:	MCTWTZJPVLRJOU-UHFFFAOYSA-N
Formula:	C4H6N2
SMILES:	Cn1ccnc1
Mol. weight [g/mol]:	82.10
CAS:	616-47-7
SMILES: Mol. weight [g/mol]: CAS:	Cn1ccnc1 82.10 616-47-7

Physical Properties

Property code	Value	Unit	Source
affp	959.60	kJ/mol	NIST Webbook
basg	927.70	kJ/mol	NIST Webbook
ie	8.66	eV	NIST Webbook
log10ws	-2.53		Crippen Method
logp	0.420		Crippen Method
mcvol	67.720	ml/mol	McGowan Method
rinpol	929.00		NIST Webbook
rinpol	929.00		NIST Webbook
rinpol	929.00		NIST Webbook
ripol	1700.00		NIST Webbook
ripol	1681.00		NIST Webbook
ripol	1638.00		NIST Webbook
tb	471.60	К	Vapor-liquid equilibrium in the production of the ionic liquid, 1-hexyl-3-methylimidazolium bromide ([HMIm][Br]), in acetone
tb	471.20	К	NIST Webbook

Temperature Dependent Properties

Property code

Unit

Source

рvар	101.33	kPa	471.60 1-he	Vapor-liquid equilibrium in the production of the ionic liquid, exyl-3-methylimidazo bromide ([HMIm][Br]), in acetone	lium
rhol	1011.90	kg/m3	323.15	Towards understanding the effect of electrostatic interactions on the density of ionic liquids	
rhol	1033.20	kg/m3	298.15	Towards understanding the effect of electrostatic interactions on the density of ionic liquids	
rhol	1029.20	kg/m3	303.15	Towards understanding the effect of electrostatic interactions on the density of ionic liquids	
rhol	1025.00	kg/m3	308.15	Towards understanding the effect of electrostatic interactions on the density of ionic liquids	
rhol	1020.70	kg/m3	313.15	Towards understanding the effect of electrostatic interactions on the density of ionic liquids	
rhol	1015.50	kg/m3	318.15	Towards understanding the effect of electrostatic interactions on the density of ionic liquids	
rhol	1037.00	kg/m3	293.15	Towards understanding the effect of electrostatic interactions on the density of ionic liquids	

rhol	1039.36	kg/m3	288.15	Mass density, sound velocity, mixing enthalpy, 1H NMR, Ab initio calculations and intermolecular interactions in binary mixtures of N-methylimidazole + water, +methanol, +ethanol, +1-propanol, +2-propanol	
rhol	1030.52	kg/m3	298.15	Mass density, sound velocity, mixing enthalpy, 1H NMR, Ab initio calculations and intermolecular interactions in binary mixtures of N-methylimidazole + water, +methanol, +ethanol, +1-propanol, +2-propanol	
rhol	1021.66	kg/m3	308.15	Mass density, sound velocity, mixing enthalpy, 1H NMR, Ab initio calculations and intermolecular interactions in binary mixtures of N-methylimidazole + water, +methanol, +ethanol, +1-propanol, +2-propanol	
rhol	1012.76	kg/m3	318.15	Mass density, sound velocity, mixing enthalpy, 1H NMR, Ab initio calculations and intermolecular interactions in binary mixtures of N-methylimidazole + water, +methanol, +ethanol, +1-propanol, +2-propanol	

rhol	1003.83	kg/m3	328.15	Mass density, sound velocity, mixing enthalpy, 1H NMR, Ab initio calculations and intermolecular interactions in binary mixtures of N-methylimidazole + water, +methanol, +ethanol, +1-propanol, +2-propanol
rhol	1031.14	kg/m3	298.15	Determination of Infinite Dilution Partial Molar Excess Enthalpies and Volumes for Some Ionic Liquid Precursors in Water and Methanol Using Tandem Flow Mixing Calorimetry and Vibrating-Tube Densimetry

Sources

Ternary Liquid-Liquid Equilibria

A combined experimental and

Influence of Typical Impurities on the Surface Tension Measurements of **Binary Mixtures of Water and the Ionic** Liquids 1-Butyl-3-Methylimidazolium Tetrafluoroborate and Chloride:

Ternary Liquid-Liquid Equilibria
Measurement for Benzene +
Cyclichekawie Equilibria Liquid: Experimental Data
routing the Liquid: Experimental Concernance
routing the https://www.doi.org/10.1021/je800376f https://www.doi.org/10.1016/j.fluid.2013.11.030 https://www.doi.org/10.1007/s10765-008-0506-x http://webbook.nist.gov/cgi/cbook.cgi?ID=C616477&Units=SI https://www.doi.org/10.1016/j.jct.2012.06.015 theoretical approach to the study of **Noticity of the study of the st** http://link.springer.com/article/10.1007/BF02311772 https://www.doi.org/10.1021/je100949x

Limiting activity coefficients of 1-chlorobutane in water and in aqueous **Sciption Metsod**stances involved in synthesis of ionic liquids: Crippen Method:

https://www.doi.org/10.1016/j.fluid.2010.09.026 https://www.chemeo.com/doc/models/crippen_log10ws http://pubs.acs.org/doi/abs/10.1021/ci990307l

Legend

affp:	Proton affinity
basg:	Gas basicity
ie:	Ionization energy
log10ws:	Log10 of Water solubility in mol/l
logp:	Octanol/Water partition coefficient
mcvol:	McGowan's characteristic volume
pvap:	Vapor pressure
rhol:	Liquid Density
rinpol:	Non-polar retention indices
ripol:	Polar retention indices
tb:	Normal Boiling Point Temperature

Latest version available from:

https://www.chemeo.com/cid/13-628-0/1H-Imidazole-1-methyl.pdf

Generated by Cheméo on 2024-04-29 02:53:04.775873103 +0000 UTC m=+16648433.696450460.

Cheméo (https://www.chemeo.com) is the biggest free database of chemical and physical data for the process industry.