

Benzoic acid, 2-(3-nitrophenyl)ethyl ester

| | |
|-----------------------------|--|
| Inchi: | InChI=1S/C15H13NO4/c17-15(13-6-2-1-3-7-13)20-10-9-12-5-4-8-14(11-12)16(18)19/h1- |
| InchiKey: | IZKRXIAUPAQEIL-UHFFFAOYSA-N |
| Formula: | C15H13NO4 |
| SMILES: | O=C(OCCc1cccc([N+](=O)[O-])c1)c1cccc1 |
| Mol. weight [g/mol]: | 271.27 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|---------|----------------------|----------------|
| gf | 92.24 | kJ/mol | Joback Method |
| hf | -146.90 | kJ/mol | Joback Method |
| hfus | 36.45 | kJ/mol | Joback Method |
| hvap | 79.95 | kJ/mol | Joback Method |
| log10ws | -4.40 | | Crippen Method |
| logp | 2.994 | | Crippen Method |
| mcvol | 199.550 | ml/mol | McGowan Method |
| pc | 2621.78 | kPa | Joback Method |
| rinpola | 2386.00 | | NIST Webbook |
| tb | 829.07 | K | Joback Method |
| tc | 1084.98 | K | Joback Method |
| tf | 539.94 | K | Joback Method |
| vc | 0.765 | m ³ /kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|--------|---------|-----------------|---------------|
| cpg | 564.32 | J/molxK | 829.07 | Joback Method |
| cpg | 576.75 | J/molxK | 871.72 | Joback Method |
| cpg | 587.93 | J/molxK | 914.37 | Joback Method |
| cpg | 597.92 | J/molxK | 957.02 | Joback Method |
| cpg | 606.79 | J/molxK | 999.67 | Joback Method |
| cpg | 614.61 | J/molxK | 1042.33 | Joback Method |
| cpg | 621.45 | J/molxK | 1084.98 | Joback Method |

Sources

| | |
|------------------------|---|
| Crippen Method: | http://pubs.acs.org/doi/abs/10.1021/ci9903071 |
| Crippen Method: | https://www.chemeo.com/doc/models/crippen_log10ws |
| Joback Method: | https://en.wikipedia.org/wiki/Joback_method |
| McGowan Method: | http://link.springer.com/article/10.1007/BF02311772 |
| NIST Webbook: | http://webbook.nist.gov/cgi/cbook.cgi?ID=U368224&Units=SI |

Legend

| | |
|-----------------|---|
| cpg: | Ideal gas heat capacity |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| hvap: | Enthalpy of vaporization at standard conditions |
| log10ws: | Log10 of Water solubility in mol/l |
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| rinpol: | Non-polar retention indices |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.chemeo.com/cid/15-560-3/Benzoic-acid-2-3-nitrophenyl-ethyl-ester.pdf>

Generated by Cheméo on 2024-05-04 16:24:05.386008704 +0000 UTC m=+17129094.306586020.

Cheméo (<https://www.chemeo.com>) is the biggest free database of chemical and physical data for the process industry.