

Butanediamide

| | |
|-----------------------------|--|
| Other names: | Succindiamide Succinic acid diamide Succinic amide butanedioic acid, diamide succinamide succinic acid, diamide succinic diamide |
| Inchi: | InChI=1S/C4H8N2O2/c5-3(7)1-2-4(6)8/h1-2H2,(H2,5,7)(H2,6,8) |
| InchiKey: | SNCZNSNPXMPCGN-UHFFFAOYSA-N |
| Formula: | C4H8N2O2 |
| SMILES: | NC(=O)CCC(N)=O |
| Mol. weight [g/mol]: | 116.12 |
| CAS: | 110-14-5 |

Physical Properties

| Property code | Value | Unit | Source |
|---------------|-----------------|---------|---|
| affp | 942.00 | kJ/mol | NIST Webbook |
| chs | -2136.10 ± 2.10 | kJ/mol | NIST Webbook |
| ep | -19.00 | J/molxK | NIST Webbook |
| gf | -142.14 | kJ/mol | Joback Method |
| hf | -283.47 | kJ/mol | Joback Method |
| hfs | -581.20 ± 2.10 | kJ/mol | NIST Webbook |
| hfus | 46.00 | kJ/mol | Odd even effect in melting properties of 12 alkane-a,x-diamides |
| hvap | 59.27 | kJ/mol | Joback Method |
| log10ws | 0.08 | | Crippen Method |
| logp | -1.263 | | Crippen Method |
| mcvol | 90.320 | ml/mol | McGowan Method |
| pc | 5390.71 | kPa | Joback Method |
| tb | 543.72 | K | Joback Method |
| tc | 761.15 | K | Joback Method |
| tf | 401.22 | K | Joback Method |
| vc | 0.330 | m3/kmol | Joback Method |

Temperature Dependent Properties

| Property code | Value | Unit | Temperature [K] | Source |
|---------------|--------|---------|-----------------|---------------|
| cpg | 235.78 | J/mol×K | 688.67 | Joback Method |
| cpg | 241.88 | J/mol×K | 724.91 | Joback Method |
| cpg | 207.08 | J/mol×K | 543.72 | Joback Method |
| cpg | 214.93 | J/mol×K | 579.96 | Joback Method |
| cpg | 222.32 | J/mol×K | 616.20 | Joback Method |
| cpg | 229.27 | J/mol×K | 652.43 | Joback Method |
| cpg | 247.57 | J/mol×K | 761.15 | Joback Method |
| cps | 174.00 | J/mol×K | 323.00 | NIST Webbook |
| hfust | 6.08 | kJ/mol | 485.90 | NIST Webbook |

Sources

Odd even effect in melting properties of <https://www.doi.org/10.1016/j.jct.2006.04.004>

12 alkane-*a,x*-diamides:
Joback Method:

https://en.wikipedia.org/wiki/Joback_method

McGowan Method:

<http://link.springer.com/article/10.1007/BF02311772>

NIST Webbook:

<http://webbook.nist.gov/cgi/cbook.cgi?ID=C110145&Units=SI>

Crippen Method:

<http://pubs.acs.org/doi/abs/10.1021/ci990307l>

Crippen Method:

https://www.chemeo.com/doc/models/crippen_log10ws

Legend

| | |
|-----------------|--|
| affp: | Proton affinity |
| chs: | Standard solid enthalpy of combustion |
| cpg: | Ideal gas heat capacity |
| cps: | Solid phase heat capacity |
| ep: | Protonation entropy at 298K |
| gf: | Standard Gibbs free energy of formation |
| hf: | Enthalpy of formation at standard conditions |
| hfs: | Solid phase enthalpy of formation at standard conditions |
| hfus: | Enthalpy of fusion at standard conditions |
| hfust: | Enthalpy of fusion at a given temperature |
| hvp: | Enthalpy of vaporization at standard conditions |
| log10ws: | Log10 of Water solubility in mol/l |

| | |
|---------------|-------------------------------------|
| logp: | Octanol/Water partition coefficient |
| mcvol: | McGowan's characteristic volume |
| pc: | Critical Pressure |
| tb: | Normal Boiling Point Temperature |
| tc: | Critical Temperature |
| tf: | Normal melting (fusion) point |
| vc: | Critical Volume |

Latest version available from:

<https://www.cheméo.com/cid/26-094-9/Butanediamide.pdf>

Generated by Cheméo on 2024-04-23 21:19:21.586055893 +0000 UTC m=+16196410.506633208.

Cheméo (<https://www.cheméo.com>) is the biggest free database of chemical and physical data for the process industry.