

# Cyclohexylamine

<b>Other names:</b>	1-Aminocyclohexane 1-Cyclohexylamine Aminocyclohexane Aminohexahydrobenzene Aniline, hexahydro- Benzenamine, hexahydro- CHA Cyclohexanamine Hexahydroaniline Hexahydrobenzenamine UN 2357 cyclohexaneamine
<b>Inchi:</b>	InChI=1S/C6H13N/c7-6-4-2-1-3-5-6/h6H,1-5,7H2
<b>InchiKey:</b>	PAFZNILMFXTMIY-UHFFFAOYSA-N
<b>Formula:</b>	C6H13N
<b>SMILES:</b>	NC1CCCCC1
<b>Mol. weight [g/mol]:</b>	99.17
<b>CAS:</b>	108-91-8

## Physical Properties

Property code	Value	Unit	Source
affp	934.40	kJ/mol	NIST Webbook
basg	899.60	kJ/mol	NIST Webbook
chl	-4071.30 ± 1.30	kJ/mol	NIST Webbook
chl	-4077.50 ± 1.60	kJ/mol	NIST Webbook
gf	90.54	kJ/mol	Joback Method
hf	-104.90 ± 1.30	kJ/mol	NIST Webbook
hf	-98.70	kJ/mol	NIST Webbook
hfl	-141.50 ± 1.60	kJ/mol	NIST Webbook
hfl	-147.70 ± 1.30	kJ/mol	NIST Webbook
hfus	8.33	kJ/mol	Joback Method
hvap	43.70	kJ/mol	NIST Webbook
hvap	42.80 ± 0.10	kJ/mol	NIST Webbook
hvap	42.80	kJ/mol	NIST Webbook
ie	8.40	eV	NIST Webbook
ie	8.40 ± 0.10	eV	NIST Webbook
ie	8.86	eV	NIST Webbook

log10ws	-1.77		Crippen Method
logp	1.278		Crippen Method
mcvol	94.520	ml/mol	McGowan Method
pc	4260.71	kPa	Joback Method
rinpol	870.00		NIST Webbook
rinpol	851.00		NIST Webbook
rinpol	857.00		NIST Webbook
rinpol	863.00		NIST Webbook
rinpol	851.00		NIST Webbook
rinpol	857.00		NIST Webbook
rinpol	855.00		NIST Webbook
rinpol	862.00		NIST Webbook
rinpol	848.00		NIST Webbook
rinpol	862.00		NIST Webbook
rinpol	857.00		NIST Webbook
ripol	1257.00		NIST Webbook
ripol	1222.00		NIST Webbook
ripol	1257.00		NIST Webbook
ripol	1265.00		NIST Webbook
ripol	1230.00		NIST Webbook
ripol	1251.00		NIST Webbook
ripol	1267.00		NIST Webbook
tb	407.00 ± 0.50	K	NIST Webbook
tb	408.15 ± 5.00	K	NIST Webbook
tb	405.15 ± 3.00	K	NIST Webbook
tb	407.70	K	NIST Webbook
tb	407.10	K	NIST Webbook
tb	407.40 ± 0.50	K	NIST Webbook
tc	626.80	K	Gas-Liquid Critical Temperatures of Some Alkenes, Amines, and Cyclic Hydrocarbons
tc	614.60	K	NIST Webbook
tf	252.00 ± 4.00	K	NIST Webbook
tf	255.40 ± 0.70	K	NIST Webbook
vc	0.334	m <sup>3</sup> /kmol	Joback Method

## Temperature Dependent Properties

Property code	Value	Unit	Temperature [K]	Source
cpg	206.73	J/mol×K	465.57	Joback Method
cpg	272.56	J/mol×K	649.65	Joback Method

cpg	260.96	J/mol×K	612.83	Joback Method
cpg	248.59	J/mol×K	576.02	Joback Method
cpg	235.45	J/mol×K	539.20	Joback Method
cpg	221.50	J/mol×K	502.39	Joback Method
cpg	191.11	J/mol×K	428.76	Joback Method
hfust	14.92	kJ/mol	255.40	NIST Webbook
hfust	16.50	kJ/mol	255.10	NIST Webbook
hvapt	40.80	kJ/mol	367.50	NIST Webbook
hvapt	40.70 ± 0.10	kJ/mol	343.00	NIST Webbook
hvapt	42.70 ± 0.10	kJ/mol	313.00	NIST Webbook
hvapt	36.14	kJ/mol	407.10	NIST Webbook
hvapt	40.80	kJ/mol	370.50	NIST Webbook
hvapt	40.60	kJ/mol	385.00	NIST Webbook
hvapt	39.60 ± 0.10	kJ/mol	358.00	NIST Webbook
pvap	11.67	kPa	343.15	An equipment for dynamic measurements of vapour liquid equilibria and results in binary systems containing cyclohexylamine
pvap	1.66	kPa	303.79	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	1.66	kPa	303.82	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures

pvap	1.66	kPa	303.84	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	2.85	kPa	313.79	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	2.85	kPa	313.81	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	4.76	kPa	323.92	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	4.76	kPa	323.97	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures

pvap	4.76	kPa	323.99	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	4.76	kPa	323.99	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	4.76	kPa	323.99	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	7.56	kPa	334.00	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	12.08	kPa	344.00	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures

pvap	17.65	kPa	354.06	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	25.96	kPa	364.10	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	25.96	kPa	364.10	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	25.94	kPa	364.13	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	25.96	kPa	364.13	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures

pvap	0.23	kPa	273.15	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	0.46	kPa	283.15	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	0.88	kPa	293.15	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	1.60	kPa	303.15	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	2.76	kPa	313.15	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures

pvap	4.57	kPa	323.15	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	7.32	kPa	333.15	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	11.35	kPa	343.15	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	17.98	kPa	353.15	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	25.06	kPa	363.15	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures



pvap	0.46	kPa	283.20	Thermodynamic properties of cyclohexanamines: Experimental and theoretical study
pvap	0.56	kPa	286.20	Thermodynamic properties of cyclohexanamines: Experimental and theoretical study
pvap	0.72	kPa	290.10	Thermodynamic properties of cyclohexanamines: Experimental and theoretical study
pvap	0.91	kPa	293.30	Thermodynamic properties of cyclohexanamines: Experimental and theoretical study
pvap	1.06	kPa	296.20	Thermodynamic properties of cyclohexanamines: Experimental and theoretical study
pvap	1.18	kPa	298.10	Thermodynamic properties of cyclohexanamines: Experimental and theoretical study
pvap	1.28	kPa	299.30	Thermodynamic properties of cyclohexanamines: Experimental and theoretical study
pvap	1.54	kPa	302.30	Thermodynamic properties of cyclohexanamines: Experimental and theoretical study
pvap	1.79	kPa	305.30	Thermodynamic properties of cyclohexanamines: Experimental and theoretical study
pvap	2.11	kPa	308.30	Thermodynamic properties of cyclohexanamines: Experimental and theoretical study

pvap	7.51	kPa	333.15	Activity Coefficients at Infinite Dilution of Cyclohexylamine + Octane, Toluene, Ethylbenzene, or Aniline and Excess Molar Volumes in Binary Mixtures of Cyclohexylamine + Heptane, Octane, Nonane, Decane, Undecane, Aniline, or Water
pvap	0.92	kPa	293.88	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	0.92	kPa	293.88	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	0.92	kPa	293.86	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures

pvap	0.92	kPa	293.83	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	0.25	kPa	274.30	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
pvap	25.59	kPa	363.15	An equipment for dynamic measurements of vapour liquid equilibria and results in binary systems containing cyclohexylamine
pvap	17.53	kPa	353.15	An equipment for dynamic measurements of vapour liquid equilibria and results in binary systems containing cyclohexylamine
pvap	7.52	kPa	333.15	An equipment for dynamic measurements of vapour liquid equilibria and results in binary systems containing cyclohexylamine

pvap	25.59	kPa	363.15	Activity Coefficients at Infinite Dilution of Cyclohexylamine + Octane, Toluene, Ethylbenzene, or Aniline and Excess Molar Volumes in Binary Mixtures of Cyclohexylamine + Heptane, Octane, Nonane, Decane, Undecane, Aniline, or Water
pvap	0.49	kPa	283.86	Experimental determination of the isothermal (vapour + liquid) equilibria of binary aqueous solutions of sec-butylamine and cyclohexylamine at several temperatures
rfi	1.46240		288.15	Vapor liquid equilibria in ternary systems of associating components (water, aniline, cyclohexylamine) and hydrocarbons (octane or toluene)
rfi	1.44599		318.15	Experimental Determination of Densities and Refractive Indices of the Ternary Mixture 2-Methyl-2-propanol + Cyclohexylamine + n-Heptane at T = (303.15 to 323.15) K
rfi	1.44856		313.15	Experimental Determination of Densities and Refractive Indices of the Ternary Mixture 2-Methyl-2-propanol + Cyclohexylamine + n-Heptane at T = (303.15 to 323.15) K

rfi	1.45114	308.15	Experimental Determination of Densities and Refractive Indices of the Ternary Mixture 2-Methyl-2-propanol + Cyclohexylamine + n-Heptane at T = (303.15 to 323.15) K
rfi	1.45375	303.15	Experimental Determination of Densities and Refractive Indices of the Ternary Mixture 2-Methyl-2-propanol + Cyclohexylamine + n-Heptane at T = (303.15 to 323.15) K
rfi	1.46240	288.15	Ternary Liquid-Liquid(-Liquid) Equilibria of Aniline + Cyclohexylamine + Water, Aniline + Cyclohexylamine + Octane, Aniline + Water + Toluene, and Aniline + Water + Octane
rfi	1.46240	288.15	Liquid-Liquid(-Liquid) Equilibria in Ternary Systems of Water + Cyclohexylamine + Aromatic Hydrocarbon (Toluene or Propylbenzene) or Aliphatic Hydrocarbon (Heptane or Octane)
rfi	1.44336	323.15	Experimental Determination of Densities and Refractive Indices of the Ternary Mixture 2-Methyl-2-propanol + Cyclohexylamine + n-Heptane at T = (303.15 to 323.15) K

rho	862.31	kg/m <sup>3</sup>	298.15	Experimental and Computational Studies of Binary Mixtures of Isobutanol + Cyclohexylamine
rho	839.55	kg/m <sup>3</sup>	323.15	Experimental determination and modelling of densities and excess molar volumes of ternary system (1-butanol + cyclohexylamine + n-heptane) and corresponding binaries from 288.15 to 323.15K
rho	844.07	kg/m <sup>3</sup>	318.15	Experimental determination and modelling of densities and excess molar volumes of ternary system (1-butanol + cyclohexylamine + n-heptane) and corresponding binaries from 288.15 to 323.15K
rho	848.61	kg/m <sup>3</sup>	313.15	Experimental determination and modelling of densities and excess molar volumes of ternary system (1-butanol + cyclohexylamine + n-heptane) and corresponding binaries from 288.15 to 323.15K
rho	857.79	kg/m <sup>3</sup>	303.15	Experimental and Computational Studies of Binary Mixtures of Isobutanol + Cyclohexylamine
rho	853.24	kg/m <sup>3</sup>	308.15	Experimental and Computational Studies of Binary Mixtures of Isobutanol + Cyclohexylamine

rho1	848.71	kg/m3	313.15	Experimental and Computational Studies of Binary Mixtures of Isobutanol + Cyclohexylamine
rho1	853.14	kg/m3	308.15	Experimental determination and modelling of densities and excess molar volumes of ternary system (1-butanol + cyclohexylamine + n-heptane) and corresponding binaries from 288.15 to 323.15K
rho1	857.67	kg/m3	303.15	Experimental determination and modelling of densities and excess molar volumes of ternary system (1-butanol + cyclohexylamine + n-heptane) and corresponding binaries from 288.15 to 323.15K
rho1	862.69	kg/m3	298.15	Thermodynamics of mixtures with strongly negative deviations from Raoult's law. XII. Densities, viscosities and refractive indices at T = (293.15 to 303.15) K for (1-heptanol, or 1-decanol + cyclohexylamine) systems. Application of the ERAS model to (1-alkanol + cyclohexylamine) mixtures

rho1	862.21	kg/m3	298.15	Experimental determination and modelling of densities and excess molar volumes of ternary system (1-butanol + cyclohexylamine + n-heptane) and corresponding binaries from 288.15 to 323.15K
rho1	866.75	kg/m3	293.15	Experimental determination and modelling of densities and excess molar volumes of ternary system (1-butanol + cyclohexylamine + n-heptane) and corresponding binaries from 288.15 to 323.15K
rho1	871.29	kg/m3	288.15	Experimental determination and modelling of densities and excess molar volumes of ternary system (1-butanol + cyclohexylamine + n-heptane) and corresponding binaries from 288.15 to 323.15K
rho1	858.12	kg/m3	303.15	Thermodynamics of mixtures with strongly negative deviations from Raoult's law. XII. Densities, viscosities and refractive indices at T = (293.15 to 303.15) K for (1-heptanol, or 1-decanol + cyclohexylamine) systems. Application of the ERAS model to (1-alkanol + cyclohexylamine) mixtures



rhoI	867.20	kg/m <sup>3</sup>	293.15	Thermodynamics of mixtures with strongly negative deviations from Raoult's law. XII. Densities, viscosities and refractive indices at T = (293.15 to 303.15) K for (1-heptanol, or 1-decanol + cyclohexylamine) systems. Application of the ERAS model to (1-alkanol + cyclohexylamine) mixtures
rhoI	866.85	kg/m <sup>3</sup>	293.15	Experimental and Computational Studies of Binary Mixtures of Isobutanol + Cyclohexylamine
speedsI	1360.60	m/s	313.15	Densities, Viscosities, Sound Speed, and IR Studies of N-methyl-2-pyrrolidone with Cyclohexylamine, Cyclohexanol, and Cyclohexene at different Temperatures.
speedsI	1342.00	m/s	318.15	Densities, Viscosities, Sound Speed, and IR Studies of N-methyl-2-pyrrolidone with Cyclohexylamine, Cyclohexanol, and Cyclohexene at different Temperatures.
speedsI	1379.20	m/s	308.15	Densities, Viscosities, Sound Speed, and IR Studies of N-methyl-2-pyrrolidone with Cyclohexylamine, Cyclohexanol, and Cyclohexene at different Temperatures.

speedsl	1301.80	m/s	323.15	Temperature dependence of densities, speeds of sound, and derived properties of cyclohexylamine + cyclohexane or benzene in the range (293.15 to 323.15) K
speedsl	1323.10	m/s	318.15	Temperature dependence of densities, speeds of sound, and derived properties of cyclohexylamine + cyclohexane or benzene in the range (293.15 to 323.15) K
speedsl	1344.50	m/s	313.15	Temperature dependence of densities, speeds of sound, and derived properties of cyclohexylamine + cyclohexane or benzene in the range (293.15 to 323.15) K
speedsl	1366.00	m/s	308.15	Temperature dependence of densities, speeds of sound, and derived properties of cyclohexylamine + cyclohexane or benzene in the range (293.15 to 323.15) K
speedsl	1408.80	m/s	298.15	Temperature dependence of densities, speeds of sound, and derived properties of cyclohexylamine + cyclohexane or benzene in the range (293.15 to 323.15) K

speedsl	1430.50	m/s	293.15	Temperature dependence of densities, speeds of sound, and derived properties of cyclohexylamine + cyclohexane or benzene in the range (293.15 to 323.15) K
speedsl	1397.80	m/s	303.15	Densities, Viscosities, Sound Speed, and IR Studies of N-methyl-2-pyrrolidone with Cyclohexylamine, Cyclohexanol, and Cyclohexene at different Temperatures.
speedsl	1387.30	m/s	303.15	Temperature dependence of densities, speeds of sound, and derived properties of cyclohexylamine + cyclohexane or benzene in the range (293.15 to 323.15) K

## Correlations

Information	Value
Property code	pvap
Equation	$\ln(P_{vp}) = A + B/T + C \cdot \ln(T) + D \cdot T^2$
Coeff. A	8.66698e+01
Coeff. B	-7.88954e+03
Coeff. C	-1.06327e+01
Coeff. D	7.33244e-06
Temperature range (K), min.	255.45
Temperature range (K), max.	615.00

## Sources

VLE and LLE in ternary systems of two associating components (water, aniline, and cyclohexylamine) and a hydrocarbon (cyclohexane or methylcyclohexane): <https://www.doi.org/10.1016/j.fluid.2014.02.023>



<b>hfust:</b>	Enthalpy of fusion at a given temperature
<b>hvap:</b>	Enthalpy of vaporization at standard conditions
<b>hvapt:</b>	Enthalpy of vaporization at a given temperature
<b>ie:</b>	Ionization energy
<b>log10ws:</b>	Log10 of Water solubility in mol/l
<b>logp:</b>	Octanol/Water partition coefficient
<b>mcvol:</b>	McGowan's characteristic volume
<b>pc:</b>	Critical Pressure
<b>pvap:</b>	Vapor pressure
<b>rfi:</b>	Refractive Index
<b>rho:</b>	Liquid Density
<b>rinpol:</b>	Non-polar retention indices
<b>ripol:</b>	Polar retention indices
<b>speedsl:</b>	Speed of sound in fluid
<b>tb:</b>	Normal Boiling Point Temperature
<b>tc:</b>	Critical Temperature
<b>tf:</b>	Normal melting (fusion) point
<b>vc:</b>	Critical Volume

Latest version available from:

<https://www.cheméo.com/cid/46-311-5/Cyclohexylamine.pdf>

Generated by Cheméo on 2024-04-23 14:59:32.946665941 +0000 UTC m=+16173621.867243252.

Cheméo (<https://www.cheméo.com>) is the biggest free database of chemical and physical data for the process industry.